



Turf Toe  
Michaud and Hauser

# THE TREATMENT AND PREVENTION OF TURF TOE: A NARRATIVE REVIEW OF CONSERVATIVE INTERVENTIONS

Tom Michaud, DC<sup>1</sup>, Andrew Hauser, MS, ATC, RSCC<sup>1</sup>

## ABSTRACT

**Objective:** This article provides an overview of the conservative management of turf toe, evaluating etiological factors associated with the development of this common injury and reviewing evidence-based treatment interventions.


**Discussion:** Turf toe is a frequently encountered sports injury in which the first metatarsophalangeal joint (MTP) is rapidly dorsiflexed, damaging some or all of the connective tissues responsible for limiting first MTP dorsiflexion. Although frequently described as a contact injury in which one player collides with the foot of another player, research shows the majority of turf toe injuries are noncontact in nature, and this distinction is significant as interventions for contact and noncontact injuries are different. The most probable biomechanical cause of noncontact turf toe is excessive pronation, as several studies have shown that this injury is more likely to occur in people with low arches. While the exact mechanism explaining the connection between turf toe and excessive pronation is unclear, this article suggests that noncontact turf toe is a continuum of functional hallux limitus, a condition in which excessive pronation increases ground reactive forces centered beneath the first metatarsal head, which in turn produces a bony blockage of first MTP dorsiflexion that greatly increases the likelihood of turf toe injury.

**Conclusion:** Specific aspects of orthotic intervention that may play a role in preventing turf toe are described in detail. The various manual techniques and

rehabilitative stretches and exercises necessary to improve first MTP motion and reduce the risk of turf toe are also discussed. (*J Contemporary Chiropr* 2026;9:207-216)

**Key Indexing Terms:** Turf Toe; Functional Hallux Limitus; Orthotic; Varus Posts; Retrotalar Pulley.

## INTRODUCTION



The term turf toe was initially introduced in 1976 by Bowers and Martin to describe a hyperextension injury of the first MTP in football players wearing flexible shoes while playing on synthetic turf. (1) The rapid and often extreme range of first MTP dorsiflexion associated with this injury is particularly likely to damage the flexor hallucis brevis tendon, the sesamoids, and the plantar plate located beneath the first metatarsal head (Fig. 1). While turf toe injuries occur in sports such as soccer, jujitsu, and rugby, they are most commonly found among professional football players, with approximately 45% experiencing this injury at some point during their careers. (2) While some authors claim that turf toe injury rates on the new synthetic turfs are comparable to turf

<sup>1</sup> Private practice of chiropractic

toe injury rates while playing on grass, a 2025 review of 1.2 million injuries over 5 years suggests otherwise. Reiad et al. (3) looked at injury rates in a wide range of sports played on different surfaces and determined compared to playing on grass, football players were 6.3 times more likely to injure their toe while playing on artificial turf. The authors cite several biomechanical studies suggesting that artificial turf exhibits greater rotational stiffness, decreased force absorption, and provides “less cleat release” than natural grass.

**Fig. 1. Anatomy of a turf toe injury.** The first MTP is reinforced by a capsuloligamentous complex, which consists of the phalangeosesamoid ligaments (PSL), the intersesamoid ligaments (ISL), the medial and lateral metatarsosesamoid ligaments (MSL), the collateral ligaments, and the plantar plate (PP). The plantar plate is a thick fibrocartilaginous part of the first MTPs joint capsule. It is reinforced medially by fibers from the abductor hallucis (AbH) and the medial head of the flexor hallucis brevis (FHB). On the lateral side, the plantar plate receives support from the oblique and transverse heads of the adductor hallucis muscle (AdH, o and t) and the lateral head of flexor hallucis brevis.

## DISCUSSION


In a frequently cited study analyzing factors contributing to the onset of turf toe, Rodeo et al. (2) reported that the primary mechanism of this injury involves a player's foot positioned on the ground with the first MTP joint dorsiflexed; subsequent contact by another player to the posterior aspect of the foot then generates a downward force, resulting in rapid and excessive dorsiflexion of the first MTP joint. This contact-related mechanism of injury is described in the vast majority of articles as the primary cause of turf toe injury in the NFL. Despite being widely referenced, turf toe injuries related to player-to-player contact are actually less frequent than previously thought. According to George et al., only 32% of these injuries involve direct contact with another player, while 35% happen without contact, typically when an athlete makes a quick change in direction, which increases both the speed and range of first MTP dorsiflexion. This is consistent with research showing that running backs and quarterbacks are more likely to experience a turf toe injury. (4)

Because the forces involved in contact-related turf toe injuries are extreme, management typically requires

increasing shoe midsole stiffness and/or using graphite insoles to limit first MTP joint dorsiflexion. In contrast, noncontact turf toe injuries may be managed by controlling first MTP joint motion during dorsiflexion and strengthening the intrinsic muscles and tendons responsible for decelerating this movement. It is important to recognize the differences between contact and noncontact turf toe injuries, as just like ACL injuries, noncontact turf toe injuries are often modifiable and amenable to specific rehab techniques that may appreciably reduce injury rates.

### *The Connection Between Turf Toe, Flat Feet, and Functional Hallux Limitus*

In a 2023 updated review of treatment options for managing turf toe, Gupta et al. (5) list “increased foot pronation” as a risk factor for developing turf toe injury. This statement is supported by Rodeo (2), who point out that among the 30 NFL players evaluated in their study following turf toe injury, nearly 75% of them possessed flat feet, which is much greater than what is typically found in the general population. There are several biomechanical reasons that people with low arches are more likely to develop turf toe. Firstly, as demonstrated by Hillstrom et al (6), excessive pronation causes the medial displacement of the talus relative to the calcaneus, which causes a medial displacement of ground reactive forces that greatly increases pressure centered beneath the hallux during propulsion. The medial displacement of force in people with low arches creates an environment in which ground reactive forces elevate the first metatarsal head relative to the second metatarsal head, engaging a bony locking mechanism that limits MTP motion. This bony locking mechanism is referred to as functional hallux limitus (Fig. 2).



**Fig. 2. Mechanism of functional hallux limitus.** In order for the first MTP to dorsiflex through a full range of motion during propulsion, the first metatarsal must actively plantarflex (A), which allows for a dorsal-posterior shifting of the transverse axis of the first MTP (B) as the metatarsal head glides posteriorly (cont.)


over the sesamoids (C). This new axis allows for an unrestrained range of hallux dorsiflexion, which improves congruency between the first metatarsal head and its proximal phalanx. When excessive foot pronation is present, medial displacement of ground reactive forces maintain the first metatarsal in an elevated position (D) thereby preventing the normal dorsal-posterior shift of the transverse axis necessary to achieve full dorsiflexion. The hallux is now forced to dorsiflex about the original axis, resulting in a decreased parallelism of the articular surfaces and a resultant jamming of the dorsal cartilage (E).

Originally described in 1972 by Laird (7) and popularized by Dananberg (8), functional hallux limitus specifically refers to a condition in which the first MTP possesses an adequate range of motion off-weight-bearing, but a limited range of motion while weight-bearing as ground reactive forces maintain the first metatarsal in an elevated position. While there are conflicting ranges cited in the literature, Halstead et al (9) claim that functional hallux limitus is present when there is greater than 50° first MTP dorsiflexion off-weight-bearing, and less than 40° of first MTP dorsiflexion while weight-bearing. Functional hallux limitus is especially common in individuals with hypermobile first rays, and this anatomical variant is frequently seen in low arched individuals.

As an individual with functional hallux limitus accelerates during propulsion, the first MTP dorsiflexes until the proximal phalanx collides with the dorsal aspect of the first metatarsal head. When this occurs, there is a superior shift in the instantaneous axis of rotation that causes the first MTP to gap open on its inferior surface (Fig. 3). Because the elevated forces placed on the first MTP are so extreme during turf toe injury, this sudden gapping creates tensile strains capable of tearing the restraining tendons, ligaments, and other soft tissues responsible for limiting movement. The abrupt restriction in first MTP dorsiflexion associated with the shifting axis is referred to as a "sagittal plane blockade," and is supported by several cadaveric studies showing that people with hallux limitus, which is extremely common in pronated foot types, have altered motion in the first MTP with dorsiflexion.

Shereff et al. (10) note that when hallux limitus is present, rather than smoothly gliding up the first metatarsal head, the proximal phalanx moves upward with a "distinctive pattern of distraction and jamming" that compresses the superior aspect of the first metatarsal head until no further motion is possible.

Limited flexibility in the plantar fascia, flexor hallucis longus (FHL), and/or flexor hallucis brevis (FHB) play a key role in the development of functional hallux limitus.



**Fig. 3. Turf toe and functional hallux limitus.** When an individual with functional hallux limitus reaches their end range of first MTP dorsiflexion, the instantaneous axis of rotation for sagittal plane motion in the first MTP shifts to the point of contact between the superior aspect of the proximal phalanx and metatarsal head (A). Because of the amplified force centered beneath the hallux in people with flat feet, the first MTP continues dorsiflexing, creating a tensile strain on the inferior aspect of the first MTP capable of producing a turf toe injury (B). The medial phalangeosesamoid ligament is typically the first ligament damaged in a turf toe injury (22), and in severe cases, both the plantar plate and stabilizing ligaments can rupture, allowing the sesamoids to retract proximally (C). Turf toe injury is classified as grade 1 if there is a stretching of the plantar complex with no ligamentous tearing and grade 2 if the plantar structures are partially torn. Grade 3 injuries involve a full tear of the plantar plate complex, causing the sesamoids to retract at least 2 mm from the proximal phalanx, which is visible on stress radiographs.

#### Orthotic Intervention


Theoretically, the biomechanical locking associated with an excessively pronated foot could be prevented if the first metatarsal was able to plantarflex. In 2006, Tom Michaud published a paper in JMPT with Deb

Nawoczenski showing that it is possible to increase first metatarsal plantarflexion during the propulsive period using a semiflexible orthotic possessing varus posts in the rearfoot and forefoot. (11) In our case study, 3D movement analysis showed the posted orthotic increased first metatarsal plantarflexion by 6° and reduced first MTP dorsiflexion by 5° while walking. Nawoczenski and Ludewig (12) had previously shown that orthotics possessing varus posts in the rearfoot and/or forefoot could increase first metatarsal plantarflexion during propulsion by an average of 3°. (9). Although one of the orthotics used in this study was posted only in the rearfoot, recent research shows that in order to control pronation during the propulsive period, which is when turf toe occurs, the best biomechanical outcomes typically occur when the orthotic is posted both in the rearfoot and forefoot, as conventional rearfoot posts alone and/or arch supports are unable to modify motion during the pushoff phase of gait. (13)

An alternate technique for modifying first MTP dorsiflexion during propulsion is to make a custom orthotic from a non-weight-bearing plaster cast taken with the first MTP fully dorsiflexed. Casting in this position causes the proximal phalanx to plantarflex the first metatarsal head, creating a positive model possessing an exaggerated decline angle to the first metatarsal. Researchers from Spain (15) used this casting technique and molded a series of 3 mm polypropylene orthotics that were made with EVA extensions beneath the forefoot with cutouts beneath the first metatarsal heads. Subsequent three-dimensional motion analysis confirmed that compared to the control group not wearing orthotics, there was nearly a 3° increase in first metatarsal plantarflexion during propulsion in subjects wearing the orthotics, which the authors claim can increase the ability of peroneus longus to stabilize the medial forefoot during propulsion. Interestingly, a prior study using this exact casting technique found that in addition to increasing first MTP motion during static stance, the individuals treated with these orthotics had significant reductions in ground reactive forces centered beneath the first metatarsal head and hallux. (16) Although these researchers did not measure MTP motion while walking or running, the redistribution of force centered beneath the medial forefoot could play an important role in decreasing the frequency of turf toe injuries.

A final consideration when using orthotics to treat/prevent turf toe injury is to bypass the orthotic shell completely and attach two-piece varus posts directly beneath the shoe's insoles. Researchers from Thailand

demonstrate the simple addition of varus wedges placed beneath the heel and forefoot of an insole significantly reduces tension in the plantar fascia during the propulsive phase of gait. (17) (Fig. 4) Because increased tension in the plantar fascia during propulsion is a driving agent for the development of functional hallux limitus as it increases compression of the proximal phalanx into the first metatarsal head, it is reasonable to assume the two-piece varus posts could reduce the potential for turf toe injury by lessening strain in the plantar fascia. In addition to lessening plantar fascial strain, these inexpensive posts reduce the overall range of rearfoot eversion and produce significant reductions in the peak range of first MTP dorsiflexion during pushoff. (17) Both factors could contribute to reducing the risk of turf toe injuries.




**Fig. 4. Two-piece varus posts placed beneath insoles.** Because increased tension in the plantar fascia (A) pulls the proximal phalanx firmly into the first metatarsal head (B), the potential for developing turf toe is greatly increased. Two-piece varus posts have been shown to reduce tension in the plantar fascia during propulsion and may therefore decrease the likelihood of subsequent turf toe injury. The thickness of the forefoot post may also be beneficial when managing turf toe injuries as it can limit the velocity of first MTP dorsiflexion during propulsion.

#### *An Alternate Cause Of Functional Hallux Limitus: Restricted Gliding of the Flexor Hallucis Longus Tendon in the Retrotalar Pulley*

A frequently overlooked cause of functional hallux limitus occurs when the flexor hallucis longus tendon gets trapped in the retrotalar pulley located behind the medial malleolus. (18) (Fig. 5) This entrapment limits excursion of the distal flexor hallucis longus tendon, which creates a compressive force on the plantar aspect of the proximal phalanx capable of blocking first MTP

motion. Restricted excursion of the FHL may also explain the unusual connection between turf toe injury and excessive ankle dorsiflexion. Originally noted by Rodeo et al. (2), athletes presenting with turf toe injury have significantly greater ranges of ankle dorsiflexion than non-injured athletes (13.3° versus 7.9°, respectively), which the authors were unable to adequately explain.


It is possible that athletes with increased ankle dorsiflexion put greater tension on their FHL tendon prior to initiating pushoff, creating a compressive force along the base of the proximal phalanx capable of engaging the first MTP locking mechanism. The rapid restriction and first MTP dorsiflexion may alter the first MTP's axis of motion, significantly increasing the risk of turf toe injury. Athletes with low arches and limited FHL excursion are especially prone to turf toe injury because of the excessive forces centered beneath the hallux in flat-footed individuals.



**Fig. 5. Entrapment of the flexor hallucis longus tendon in the retrotalar pulley.** The tendon of flexor hallucis longus passes through a small tunnel located between the posteromedial (A) and posterolateral (B) processes of the talus. Adhesions between the tendon and its synovial lining at the pulley and/or a low lying FHL muscle belly (18) can restrict gliding of the flexor hallucis longus tendon through the pulley, an effect Hamilton (36) likens to a cork becoming lodged in a bottle.

Fortunately, excursion of the FHL tendon is easily diagnosed by performing the FHL stretch test. (19) This test is performed by measuring first MTP dorsiflexion with the ankle plantarflexed slightly, and then again with the ankle fully dorsiflexed. The test is positive when first

MTP dorsiflexion is significantly limited when measured with the ankle fully dorsiflexed. Researchers from Switzerland note that it is possible to correct limited excursion of the FHL tendon with a simple manual technique known as the Hoover Cord Maneuver. (20) This maneuver is performed by grasping the heel with one hand while simultaneously dorsiflexing the forefoot with the other (Fig. 6). The hand grasping the heel creates a firm distraction force while gently rocking the calcaneus from full inversion to full eversion. According to the authors, this maneuver almost always allows for the full restoration of first MTP dorsiflexion, which is immediately apparent to the patient, "who is often surprised to see the recovery of the great toe's range of motion after the manipulation." The authors performed this mobilization on 16 participants and noted immediate increases in first MTP motion in every subject. To prevent flexor hallucis longus tendon entrapment in the retrotalar pulley, patients should perform bent-knee calf stretches with a towel under the big toe. Each stretch should be held for 30 seconds and repeated five times daily.



**Fig. 6. The Hoover Cord Maneuver.** With the patient resting supine, both hands work together to simultaneously traction subtalar joint while maintaining the ankle in a slightly dorsiflexed position (arrows). While maintaining constant traction, the hand contacting the heel gently sways the subtalar joint through a full range of inversion/eversion. According to Vallotton et al. (20), these movements encourage distal glide of the tendon through the pulley thereby increasing the length of the FHL tendon between the pulley and its insertion on the distal phalanx.


### The effect of Limited First MTP Motion Following Injury

Regardless of whether the turf toe injury was contact or noncontact related, prior injury is notorious for limiting first MTP motion. Brophy et al. (21) showed that football players with a prior history of turf toe had an 8° reduction in first MTP dorsiflexion compared to uninjured athletes (40.6 versus 48.4°, respectively). This significant limitation of motion may predispose the restraining structures to tear should the MTP be forced to rapidly dorsiflex beyond the reduced range. While first MTP motion may be limited following injury because of plantar fascial and/or FHL contracture, one of the most common causes for limited MTP motion following a dorsiflexion injury is contracture of the FHB muscle.

This is easily distinguished from impaired gliding of the FHL because with FHB contracture, first MTP motion is limited with the ankle in a neutral and a dorsiflexed position, while FHL contracture only limits MTP dorsiflexion with the ankle dorsiflexed. When FHB contracture is present, it is easy to restore dorsiflexion with the muscle energy stretch illustrated in Fig. 7.

This muscle energy stretch is also used to determine if contracture in the plantar fascia is limiting first MTP motion, as the plantar fascia is naturally stiff and inflexible, it does not lengthen in response to gentle hold relax stretches. Although occasionally problematic, contracture of the plantar fascia is a rare cause of post-traumatic limited MTP motion, as the overwhelming majority of people presenting with turf toe injuries have rapid restoration of motion with hold-relax stretching.

Although some authorities advise that athletes not return to sport until off-weight-bearing dorsiflexion reaches 50 to 60°, evidence from a cadaveric study by Framenco et al. (22) indicates that most turf toe injuries occur at angles below 78°. Consequently, it is recommended that the target for off-weight-bearing first MTP dorsiflexion on both the injured and uninjured sides be set at 75–80°. Of course, this target goal may not be possible for all athletes, but the mobilization illustrated in figure 7 produces surprisingly large increases in first MTP dorsiflexion.



**Fig. 7. Neutral-position stretch to lengthen flexor hallucis brevis.** To begin this muscle-energy stretch, the patient plantarflexes the first MTP with mild force while the practitioner maintains the hallux in a midline position (A). This light pressure is maintained for five seconds. Next, the practitioner moves the first MTP to the limit of its available range of dorsiflexion by using the thumb of one hand to displace the first metatarsal head inferiorly (B), while the opposite hand glides the proximal phalanx superiorly (C). The joint is held in this stretched position for 10 seconds. The first MTP is then moved back to its neutral position, and the entire process is repeated four times. Because the isometric contractions are performed gently with the first MTP maintained in a midline position, this stretching technique is less likely to injure the plantar plate and/or restraining ligaments than conventional muscle energy stretches.

### The Overlooked Role of Strengthening Muscles and Tendons

In order to resist the extreme forces associated with turf toe injury, it is important that the muscles and tendons that limit first MTP motion be as strong as possible. A powerful FHL and FHB can protect against excessive sagittal plane motions, while abductor and adductor hallucis can stabilize the first MTP in the transverse plane. Although it has no direct attachment to the first MTP, peroneus longus is an especially important muscle because it is the most powerful stabilizer of the medial forefoot. As demonstrated by Brophy et al (21), athletes with prior turf toe injury generate significantly less pressure beneath their first metatarsal heads compared to uninjured athletes, making peroneus longus strengthening essential for the treatment and prevention of turf toe injuries.

Needless to say, for exercises to work well, they should boost both muscle activation and force production. Surprisingly, that doesn't always happen. Osborne et al. (23) recently assessed EMG activity and torque generated at the metatarsophalangeal joints with 16 commonly prescribed foot and ankle exercises. Their findings revealed an intriguing paradox: certain exercises elicited substantial muscle activation, yet these EMG increases did not correspond to greater force output beneath the forefoot. Specifically, the short foot, squat, and toe spread exercises, which are frequently prescribed when managing turf toe injuries, were particularly ineffective at increasing metatarsophalangeal joint torque. They state the mismatch between muscle activation and force output occurs because the muscles are exercised in their shortened positions, which greatly impairs force production. Following this logic, other commonly prescribed exercises for managing turf toe, such as towel curl and marble pickup exercises, would also produce negligible changes in force output toes and should therefore be avoided.

Goldman (24) showed that optimal effectiveness in strengthening the toe muscles is achieved when exercises are performed with toe muscles in their lengthened positions, resulting in strength gains that are four times greater than those observed with conventional exercise methods. In addition to strengthening muscles, tendons also respond well to being exercised in their lengthened positions, particularly with prolonged isometric contractions. Several studies have shown that performing as few as 4, 30-second isometric contractions with tendons maintained in their lengthened positions produces appreciable gains in tendon strength (25,26), which would allow them to tolerate greater force before failure.

#### Evidence-Based Exercise Prescription

Turf toe rehabilitation is typically broken down into four stages depending upon the severity of injury. The primary goal of phase 1 exercises is to enhance foot strength and mobility without placing strain on the first metatarsophalangeal (MTP) joint (Fig. 8). In contrast, phase 2 exercises introduce controlled motion of the first MTP joint within a limited range, applying minimal force. Elastic bands are used for most phase 2 exercises, and the timing of movement is set to the beat of metronome, which has been proven to reduce intracortical inhibition and increase motor output following injury (27). For phase 1 and 2 exercises, a regimen consisting of four sets of 25 repetitions performed to fatigue is commonly recommended, as

this protocol has been shown to promote significant muscle hypertrophy even with low resistance. (28) After completing each set, athletes are instructed to perform a 30-second isometric contraction at 60 to 90% of maximal effort, maintaining the muscles in a lengthened position throughout. Perhaps the most important phase 2 exercise is the partial forward lean maneuver, as this simple exercise has been shown to increase force output at the metatarsophalangeal joint by 78%. (23) Because the force centered beneath the hallux with this exercise is so high, early stages of this exercise are performed by leaning forward just a small amount, gradually increasing the degree of forward leaning as strength improves.





Fig. 8. Phase 1 and 2 turf toe exercises.

As athletes progress to phase 3 (Fig. 9), elastic band exercises are replaced with more aggressive weight-bearing exercises, which are performed with muscles in their lengthened positions. It is possible to determine if an athlete is ready to begin phase 3 exercises by measuring force generated beneath the hallux with a toe strength dynamometer. This device has excellent interrater reliability (31) and athletes should be able to generate 10% body weight beneath their hallux before beginning these exercises. One of the most effective phase 3 exercise is the ToePro platform. A pilot study from Temple University (29) demonstrated that performing ToePro exercises three times a week for six weeks led to a 35% increase in force generated beneath the hallux, with improvements in single-leg balance. The peroneal/FHL exercise ToePro exercise is particularly effective at increasing force output beneath the medial forefoot. As the athlete gets stronger, the intensity of these exercises gradually increases and as strength increases, prolonged isometric contractions are added

at the end of each set, which in addition to improving tendon strength, also act as a natural analgesic. (30)



**Fig. 9. Phase 2 and 3 turf toe exercises.** A comprehensive written review and an accompanying video covering the complete turf toe rehabilitation program, including recommended sets, repetitions, intensities, hold times and progressions is available at [www.humanlocomotion.com](http://www.humanlocomotion.com).

The phase 4 exercises make the transition away from low to moderate load/high repetition exercise routines towards heavy load/low repetition exercises. Many of the phase 4 exercises were adapted from an article by Tourillon et al. (35), who performed an 8-week strength training routine on 28 highly-trained athletes. They discovered that these exercises not only strengthened the muscles that stabilize the first MTP joint, which is crucial for preventing turf toe, they also improved athletic performance. Specifically, participants demonstrated enhanced cutting ability, better side-to-side force transfer, and greater vertical propulsion at top speeds.

These benefits are valuable for athletes at any skill level. Although not illustrated, the final stage 4 exercises include various plyometric drills, such as single leg hops, aggressive sprinting with changes in direction, four-way hip strengthening, and a variety of core exercises. Because these drills are aggressive, athletes should be monitored closely.

## CONCLUSION

Comparable to ACL injuries, turf toe should be classified as being either a contact or noncontact related injury. While contact related turf toe injuries may be managed with continued improvements in synthetic turf technology and changes in shoe gear, noncontact related injury should be managed with specific interventions that correct the faulty first MTP motions associated with functional hallux limitus; i.e., pronators should be treated with orthotics or insole modifications possessing varus posts in both the forefoot and the rearfoot, and individuals with either FHB or FHL contracture should be identified and treated with the appropriate manual techniques.

Although there is no current research directly connecting functional hallux limitus to turf toe injury, the recommended interventions are safe with little potential to produce injury. Advanced strengthening exercises should be encouraged, as multiple studies have demonstrated that foot strengthening routines not only reduce the overall incidence of injuries among athletes (32), but also result in notable improvements in speed, jumping ability, and agility. (24,33-35)

## REFERENCES

1. Bowers Jr KD, Martin RB. Turf-toe: a shoe-surface related football injury. *Med Sci Sports*. 1976;8(2):81-83
2. Rodeo SA, O'Brien S, Warren RF, Barnes R, Wickiewicz TL, Dillingham MF. Turf-toe: an analysis of metatarsophalangeal joint sprains in professional football players. *AM J Sports Med*. 1990;18(3):280-285
3. Reiad TA, Milner JD, Dinh PV et al. Comparing sports-related orthopedic injury trends on artificial turf and natural grass: a 20-year nationwide analysis of the NEISS Database. *Rhode Island Med J*. 2025;108(7)
4. George E, Harris AH, Dragoo JL, Hunt KJ. Incidence and risk factors for turf toe injuries in intercollegiate football: data from the national collegiate athletic association injury surveillance system. *Foot Ankle International*. 2014;35(2):108-115
5. Gupta A, Singh PK, Xu AL, Bronheim RS, McDaniel CM, Aiyer AA. Turf toe injuries in the athlete: an updated review of treatment options, rehabilitation

- protocols, and return-to-play outcomes. *Current Rev Musculoskeletal Med.* 2023;16(11):563-574
6. Hillstrom HJ, Song J, Kraszewski AP et al. Foot type biomechanics part 1: structure and function of the asymptomatic foot. *Gait Posture.* 2013;37(3):445-451
  7. Laird PO. Functional hallux limitus. *Illinois Podiatrists.* 1972;9(4)
  8. Dananberg HJ. Gait style as an etiology to chronic postural pain. Part I. Functional hallux limitus. *J Am Podiatric Med Assoc.* 1993;83(8):433-441
  9. Halstead J, Redmond A. Weight-bearing passive dorsiflexion of the hallux in standing is not related to hallux dorsiflexion during walking. *J Orthop Sports Phys Ther.* 2006;36:550.
  10. Shereff MJ, Bejjani FJ, Kummer FJ. Kinematics of the first metatarsophalangeal joint. *JBJS.* 1986;68(3):392-398
  11. Michaud TC, Nawoczenski DA. The influence of two different types of foot orthoses on first metatarsophalangeal joint kinematics during gait in a single subject. *J Manipulative Physiol Ther.* 2006;29(1):60-65
  12. Nawoczenski DA, Ludewig PM. The effect of forefoot and arch posting orthotic designs on first metatarsophalangeal joint kinematics during gait. *J Orthop Sports Phys Ther.* 2004;34(6):317-327
  13. Jor A, Lau NW, He Y et al. Effects of foot orthoses on lower extremity joint kinematics and kinetics in runners with asymptomatic flatfeet: A systematic review and meta-analysis. *Gait Posture.* 2025 Jun 6
  14. Welsh BJ, Redmond AC, Chockalingam N, Keenan AM. A case-series study to explore the efficacy of foot orthoses in treating first metatarsophalangeal joint pain. *J Foot Ankle Res.* 2010;3(1):17
  15. Becerro de Bengoa Vallejo R, Gómez RS, Losa Iglesias ME. Clinical improvement in functional hallux limitus using a cut-out orthosis. *Prosthetics Orthotics International.* 2016;40(2):215-223
  16. Scherer PR, Sanders J, Eldredge DE, Duffy SJ, Lee RY. Effect of functional foot orthoses on first metatarsophalangeal joint dorsiflexion in stance and gait. *J Am Podiatric Med Assoc.* 2006;96(6):474-481
  17. Harutaichun P, Vongsirinavarat M, Pakpakorn P, Sathianpantarit P, Richards J. Can orthotic wedges change the lower-extremity and multi-segment foot kinematics during gait in people with plantar fasciitis?. *Gait Posture.* 2022;97:174-183
  18. Sockalingam N, Reymond N, Rybnikov A, Dubois-Ferriere V, Assal M. Do patients with functional hallux limitus have a low-lying or bulky fhl muscle belly? *Foot Ankle Orthop.* 2023;8(1):24730114231153140
  19. Michelson J, Dunn L. Tenosynovitis of the flexor hallucis longus: a clinical study of the spectrum of presentation and treatment. *Foot Ankle International.* 2005;26(4):291-303
  20. Vallotton J, Echeverri S, Dobbelaere-Nicolas V. Functional hallux limitus or rigidus caused by a tenodesis effect at the retrotalar pulley: Description of the functional stretch test and the simple hoover cord maneuver that releases this tenodesis. *J Am Podiatric Med Assoc.* 2010;100(3):220-229
  21. Brophy RH, Gamradt SC, Ellis SJ et al. Effect of turf toe on foot contact pressures in professional American football players. *Foot Ankle International.* 2009;30(5):405-409
  22. Frimenko RE, Lievers WB, Riley PO et al. Development of an injury risk function for first metatarsophalangeal joint sprains. *Med Sci Sports Exercise.* 2013 Nov 1;45(11):2144-2150
  23. Osborne JW, Menz HB, Landorf KB, Whittaker GA, Cotchett M, Kelly LA. The influence of body posture and added mass on intrinsic and extrinsic foot muscle activation and force output during common foot strengthening exercises. *J Sport Health Sci.* 2025:101110
  24. Goldmann JP, Sanno M, Willwacher S, Heinrich K, Brüggemann GP. The potential of toe flexor muscles to enhance performance. *J Sports Sci.* 2013;31(4):424-433
  25. Paxton JZ, Hagerty P, Andrick JJ, Baar K. Optimizing an intermittent stretch paradigm using ERK1/2 phosphorylation results in increased collagen synthesis in engineered ligaments. *Tissue engineering part A.* 2012;18(3-4):277-284
  26. Oranchuk DJ, Storey AG, Nelson AR, Cronin JB. Isometric training and long-term adaptations: Effects of muscle length, intensity, and intent: A systematic review. *Scandinavian J Med Sci Sports.* 2019;29(4):484-503

27. Gordon T, Jeanfavre M, Leff G. Effects of tempo-controlled resistance training on corticospinal tract plasticity in healthy controls: a systematic review. *Healthcare* 2024;12:1325).
28. Burd NA, West DW, Staples AW et al. Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One*. 2010;5(8):e12033
29. Song J, Gorelik S, Husang D, Morgan T. Effects of eccentric exercises on foot structure, balance, and dynamic plantar loading. Gait Study Center, Temple University School of Podiatric Med. 2019.
30. Rio E, Kidgell D, Purdam C, Gaida J, Moseley GL, Pearce AJ, Cook J. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med*. 2015;49(19):1277-1283
31. Xu J, Goss DD, Saliba SA. A novel intrinsic foot muscle strength dynamometer demonstrates moderate-to-excellent reliability and validity. *International J Sports Phys Ther*. 2023;18(4):997
32. Taddei UT, Matias AB, Duarte M, Sacco IC. Foot core training to prevent running-related injuries: a survival analysis of a single-blind, randomized controlled trial. *Am J Sports Med*. 2020;48(14):3610-3619.
33. Sulowska I, Mika A, Oleksy Ł, Stolarczyk A. The influence of plantar short foot muscle exercises on the lower extremity muscle strength and power in proximal segments of the kinematic chain in long-distance runners. *BioMed Res International*. 2019;2019(1):6947273
34. Yuasa Y, Kurihara T, Isaka T. Relationship between toe muscular strength and the ability to change direction in athletes. *J Human Kinetics*. 2018;64:47
35. Tourillon R, Fourchet F, Edouard P, Morin JB. Effects of a forefoot strengthening protocol on explosive tasks performance and propulsion kinetics in athletes: a single-blind randomized controlled trial. *PLoS One*. 2025;20(6):e0313979.
36. Hamilton WG. Stenosing tenosynovitis of the flexor hallucis longus tendon and posterior impingement upon the os trigonum in ballet dancers. *Foot Ankle*. 1982;3(2):74-80